arXiv:1305.1190v1 [math.CO] 6 May 2013

Cubic symmetric graphs having an abelian automorphism group with two orbits

Hiroki Koike
University of Primorska, UP IAM
Muzejski trg 2, SI6000 Koper, Slovenia
`hiroki.koike@upr.si`

István Kovács
University of Primorska, UP IAM and UP FAMNIT
Muzejski trg 2, SI6000 Koper, Slovenia
`istvan.kovacs@upr.si`

May 7, 2013

Abstract

Finite connected cubic symmetric graphs of girth 6 have been classified by K. Kutnar and D. Marušič (*J. Combin. Theory Ser. B* **99** (2009), 162–184), in particular, each of these graphs has an abelian automorphism group with two orbits on the vertex set. In this paper all cubic symmetric graphs with the latter property are determined. In particular, with the exception of the graphs $K_4, K_{3,3}, Q_3, GP(5,2), GP(10,2), F40$ and $GP(24,5)$, all the obtained graphs are of girth 6.

Keywords: cubic symmetric graph, Haar graph, voltage graph.

MSC 2010: 20B25, 05C25.

1 Introduction

In this paper all graphs are finite, simple and connected. For a graph X, we let $V(X)$, $E(X)$, $A(X)$, and $\mathrm{Aut}(X)$ denote the vertex set, the edge set, the arc set, and the full group of automorphisms of X, respectively. For an integer $s \geq 1$, by an *s-arc* of X we mean an ordered $(s+1)$-tuple $(v_0, \ldots, v_s)$ of vertices such that $v_i \sim v_{i+1}$ for every $i \in \{0, \ldots, s-1\}$, and if $s \geq 2$, then $v_i \neq v_{i+2}$ for every $i \in \{0, \ldots, s-2\}$. A graph X is called *s-regular* when $\mathrm{Aut}(X)$ is regular on the set of its s-arcs, and it is called *symmetric* when the group $\mathrm{Aut}(X)$ is transitive on $A(X)$.

This paper concerns with cubic symmetric graphs, i.e., connected regular graphs X of degree 3, for which $\mathrm{Aut}(X)$ is transitive on $A(X)$. It was proved by Tutte [20] that every such graph is s-regular for some $s \leq 5$. This result was improved by Djoković and Miller [8], who classified arc-transitive group actions on symmetric cubic graphs into seven types. For a recent development on this classification, we refer to [6]. In studying cubic symmetric graphs, many authors considered those graphs that are, in addition, of girth 6 (see, e.g. [5, 10, 15, 17, 18, 19]). The classification of these graphs has been completed by Kutnar and Marušič [15]. It follows that, if X is a cubic symmetric graph of girth 6, then $\mathrm{Aut}(X)$ contains an abelian subgroup with two orbits on $V(X)$ (see [15, Proposition 2.1, Theorems 1.1 and 1.2]). This observation motivated the present paper, where the object, we have proposed to ourself, is to find all cubic symmetric graphs having an abelian automorphism group with two orbits on the vertex set. It is worth to note that, the class of graphs considered by us includes all Cayley graphs on a group which has an abelian subgroup of index 2; the latter graphs have been the subject in a few papers (see, e.g. [1, 7, 22]).

Before stating the main result we set the necessary notations: $Q_3, K_n, K_{n,n}, \mathrm{F}n$ denote the cube, the complete graph on n vertices, the complete bipartite graph with color classes each of size n, and the cubic symmetric graph on n vertices provided that this is uniquely determined by n. The graphs described below will give the bulk of our classification:

Example 1.1. (Generalized Petersen graphs) Let n, k be positive integers, $n \geq 3$, and $n/2 > k$, then the *generalized Petersen graph* $GP(n,k)$ has vertex set $\{u_i, v_i : i \in \{0, ..., n-1\}\}$, and its edges are in the form:

$$\{u_i, u_{i+1}\}, \{u_i, v_i\} \text{ and } \{v_i, v_{i+k}\},$$

where the arithmetic in the indices is done modulo n. Clearly, $GP(n,k)$ is a connected cubic graph. The mapping $u_i \mapsto u_{i+1}, v_i \mapsto v_{i+1}$ is an automorphism, and this generates a group with two orbits on the vertex set. The following result is [11, Theorem A]:

Frucht-Graver-Watkins Theorem (1971). *The graph $GP(n,k)$ is symmetric if and only if $(n,k) \in \{(4,1), (5,2), (8,3), (10,2), (10,3), (12,5), (24,5)\}$.*

To sum up, $GP(4,1), GP(5,2), GP(8,3), GP(10,2), GP(10,3), GP(12,5)$ and $GP(24,5)$ are connected cubic symmetric graphs having an abelian (in fact, cyclic) automorphism group with two orbits on the vertex set. □

Example 1.2. (Haar graphs) Let G be a finite group, a, b be in G such that $a \neq b$ and $G = \langle a, b \rangle$. Then the *Haar graph* $H(G; a, b)$ has vertex set $\{u_g, v_g : g \in G\}$, and its edges are in the form:

$$\{u_g, v_g\}, \{u_g, v_{ga}\} \text{ and } \{u_g, v_{gb}\}.$$

Haar graphs were introduced in [12], and were redefined under the name *bi-Cayley graphs* in [21]. Clearly, $H(G; a, b)$ is connected and cubic. Moreover, the action of G

on the vertex set, given by $u_x \mapsto u_{xg}, v_x \mapsto v_{xg}$, induces an automorphism group with two orbits; the latter group will be denoted also by G. Suppose, in addition, that G is abelian, and there exists some $\sigma \in \mathrm{Aut}(G)$ such that $a^\sigma = ba^{-1}$ and $b^\sigma = a^{-1}$. Set $X = H(G; a, b)$, and define the mappings α and β of $V(X)$ by

$$\alpha : u_g \mapsto u_{g^\sigma},\ v_g \mapsto v_{g^\sigma a} \text{ and } \beta : u_g \mapsto v_{g^{-1}},\ v_g \mapsto u_{g^{-1}}.$$

A direct check shows that both α and β normalize G, and both are automorphisms of X. The group $\langle G, \alpha, \beta\rangle$ is transitive on $V(X)$, and α fixes the vertex u_{1_G} and permutes the neighbors of u_{1_G} in the 3-cycle $(v_{1_G}\, v_a\, v_b)$. Here and in what follows 1_G will denote the identity element of G. These imply that $\langle G, \alpha, \beta\rangle$ is transitive on $A(X)$, and therefore, $X = H(G; a, b)$ is a symmetric cubic graph. □

The main result of this paper is the following theorem:

Theorem 1.3. *Let X be a finite connected cubic symmetric graph, and $G \leq \mathrm{Aut}(X)$ be an abelian subgroup with two orbits on $V(X)$. Then one of the following two cases holds:*

(A) G acts unfaithfully on one of its orbits, and (X, G) is isomorphic to one of the following pairs:

$(K_4, \mathbb{Z}_3), (K_4, \mathbb{Z}_2^2), (K_{3,3}, \mathbb{Z}_3^2), (K_{3,3}, \mathbb{Z}_2^2), (K_{3,3}, \mathbb{Z}_4), (Q_3, \mathbb{Z}_6), (Q_3, \mathbb{Z}_2^3), (GP(8,3), \mathbb{Z}_{12})$.

(B) G acts faithfully on both of its orbits, and either $X \cong H(G; a, b), G = \langle a, b\rangle, a^\sigma = ba^{-1}$ and $b^\sigma = a^{-1}$ for some $\sigma \in \mathrm{Aut}(G)$; or (X, G) is isomorphic to one of the following pairs:

$(K_4, \mathbb{Z}_2), (Q_3, \mathbb{Z}_4), (Q_3, \mathbb{Z}_2^2), (GP(5,2), \mathbb{Z}_5), (GP(8,3), \mathbb{Z}_8), (GP(8,3), \mathbb{Z}_2 \times \mathbb{Z}_4)$,

$(GP(10,2), \mathbb{Z}_{10}), (GP(10,3), \mathbb{Z}_{10}), (GP(12,5), \mathbb{Z}_{12}), (GP(12,5), \mathbb{Z}_2 \times \mathbb{Z}_6), (GP(24,5), \mathbb{Z}_{24})$,

$(F40, \mathbb{Z}_2 \times \mathbb{Z}_{10})$.

Remark 1.4. We remark that, with the exceptions of the following seven graphs:

$$K_4,\ K_{3,3},\ Q_3,\ GP(5,2),\ GP(10,2),\ F40,\ GP(24,5),$$

every graph in Theorem 1.3 is of girth 6. To see this, we observe first that the Haar graph $H(G; a, b)$ with an abelian group G has the 6-circuit:

$$v \sim u \sim va \sim uab^{-1} \sim vab^{-1} \sim ub^{-1} \sim v,$$

and hence the girth of $H(G; a, b)$ is 4 or 6 (the graph is bipartite). Actually, the girth is 6, unless $H(G; a, b) \cong K_{3,3}$ or Q_3 (it is well-known that K_4, $K_{3,3}$ and Q_3 are the only connected cubic symmetric graphs of girth less then 5). Then we obtain the above seven graphs, after we have checked, simply using [4, Table], the girth of the remaining graphs in Theorem 1.3. In this paper we shall occasionally need information about cubic symmetric graphs of small order, and for this purpose shall use the catalog [4, Table].

We prove part (A) of Theorem 1.3 in Section 2, and part (B) in Sections 3–4. For the rest of the paper we keep the following notations:

- X is a connected cubic symmetric graph.
- $G \leq \mathrm{Aut}(X)$ is an abelian group with two orbits on $V(X)$.
- U and V are the orbits of G on $V(X)$.

2 Proof of part (A)

In this section we consider the case when G is unfaithful on U or V. We start with a simple lemma which will be used in the proof of Theorem 1.3.(A). Given a circuit C of X, we denote by $E(C)$ the set of edges which appear in C.

Lemma 2.1. *With notations as above, let F be a subset of $E(X)$, and $\mathcal{C}$ be the set of all ℓ-circuits of X for some $\ell \geq 2$. If $\mathcal{C}$ is not empty, then*

$$\frac{1}{|\mathcal{C}|}\sum_{C\in\mathcal{C}} |F \cap E(C)| = \frac{\ell \cdot |F|}{|E(X)|}. \tag{1}$$

Proof. Since $\mathrm{Aut}(X)$ is transitive on the set $E(X)$, the number of ℓ-circuits on a given edge $e \in E(X)$ does not depend on the choice of e; let this number be denoted by k. Let us count the pairs $(e, C) \in E(X) \times \mathcal{C}$ for which $e \in E(C)$. The standard double-counting gives

$$k \cdot |E(X)| = \ell \cdot |\mathcal{C}|.$$

Now, count those pairs $(e, C) \in F \times \mathcal{C}$ for which $e \in E(C)$. Then

$$\sum_{C\in\mathcal{C}} |F \cap E(C)| = k \cdot |F| = \frac{\ell \cdot |\mathcal{C}| \cdot |F|}{|E(X)|},$$

and formula (1) holds. □

It is well-known that if X has girth at most 4, then $X \cong K_4$ or $K_{3,3}$ or Q_3. These graphs have several abelian automorphism groups with two orbits, and since these groups will appear throughout the paper, we have recorded them, up to conjugacy, in Table 1. In this table N denotes the largest normal subgroup of $\mathrm{Aut}(X)$ contained in G, and $\mathbf{m}$ denotes the number of complete matchings between the two orbits of G provided that these are of the same length.

Proof of Theorem 1.3.(A). If X has girth at most 4, then the first seven pairs in Theorem 1.3.(A) follow by Table 1. For the rest of the proof let the girth of X be larger than 4. Suppose that G acts unfaithfully on U, and let G_U be the kernel of G acting on U, i.e., $G_U = \{g \in G : ug = u \text{ for every } u \in U\}$. Let $g \in G_U$ and $v \in V$ such that $vg \neq v$. If $v \sim u_1$ and $v \sim u_2$ for distinct vertices $u_1, u_2 \in U$, then $vg \sim u_ig = u_i$ for $i = 1, 2$, and hence $v \sim u_1 \sim vg \sim u_2 \sim v$. This is impossible as the girth is larger

X	G	N	faithful	degrees	**m**	X	G	N	faithful	degrees	**m**
K_4	$\mathbb{Z}_3$	1	NO	$1,3$	–	Q_3	$\mathbb{Z}_6$	$\mathbb{Z}_2$	NO	$2,4$	–
K_4	$\mathbb{Z}_2^2$	$\mathbb{Z}_2^2$	NO	$2,2$	2	Q_3	$\mathbb{Z}_2^3$	$\mathbb{Z}_2$	NO	$4,4$	2
K_4	$\mathbb{Z}_2$	1	YES	$2,2$	2	Q_3	$\mathbb{Z}_4$	1	YES	$2,2$	1
						Q_3	$\mathbb{Z}_2^2$	1	YES	$2,2$	1
$K_{3,3}$	$\mathbb{Z}_4$	1	NO	$2,4$	–	Q_3	$\mathbb{Z}_2^2$	1	YES	$4,4$	2
$K_{3,3}$	$\mathbb{Z}_2^2$	1	NO	$2,4$	–	Q_3	$\mathbb{Z}_2^2$	$\mathbb{Z}_2$	YES	$4,4$	2
$K_{3,3}$	$\mathbb{Z}_3^2$	1	NO	$3,3$	3	Q_3	$\mathbb{Z}_4$	1	YES	$4,4$	3
$K_{3,3}$	$\mathbb{Z}_3$	1	YES	$3,3$	3	Q_3	$\mathbb{Z}_2^2$	$\mathbb{Z}_2^2$	YES	$4,4$	3

Table 1: Abelian subgroups $G \leq \mathrm{Aut}(X)$ with two orbits on $V(X)$.

than 4. Using also that U, V are orbits of G and that X is connected, this gives that every vertex in V has exactly one neighbor in U. Now, if G acts unfaithfully also on V, then the same argument yields that any vertex in U has exactly one neighbor in V. Therefore, if $g \in G_U$, then g fixes pointwise V. This, however, contradicts that G_U is non-trivial. We conclude that G acts faithfully on V. Fix a vertex $v \in V$. Since G is faithful on V, it is regular on V, and we may write $V = \{vg : g \in G\}$, where $vg \neq vg'$ whenever $g \neq g'$.

Let $X[V]$ denote the subgraph of X induced by V. This graph is regular of degree 2, hence there exists $a \in G$ such that

$$E(X[V]) = \big\{\{vg, vga\} : g \in G\big\}.$$

As the girth is at least 5, we see that the order of a is at least 5. In what follows the order of a will be denoted by $|a|$.

Since G is transitive on V and commutes with G_U, G_U is semiregular on V. Then every orbit of G_U has length $|G_U|$. For a vertex $vg \in V$, denote by $\mathrm{Orb}_{G_U}(vg)$ the orbit of vg under G_U. There is a unique vertex $u \in U$ such that $u \sim w$ for every $w \in \mathrm{Orb}_{G_U}(vg)$. Thus $|G_U| = |\,\mathrm{Orb}_{G_U}(vg)| \leq 3$, and so $|G_U| = 2$ or 3. We set b for a generator of G_U. Since $v \sim va$, $\mathrm{Orb}_{G_U}(v) \neq \mathrm{Orb}_{G_U}(va)$. Thus there exist $u_1, u_2 \in U, u_1 \neq u_2$, such that we have the 6-circuit:

$$C : v \ \sim \ u_1 \ \sim \ vb \ \sim \ vba \ \sim \ u_2 \ \sim \ va \ \sim \ v. \tag{2}$$

Case 1. $|G_U| = 3$.

Since no edge lies in U and X is connected, the group $\langle a, b\rangle$ is transitive on V. Using also that G is regular on V, we obtain that $G = \langle a, b\rangle$. Also, $|V| = |G|$, $|U| = \frac{1}{3}|G|$, $|V(X)| = \frac{4}{3}|G|$, and $|E(X)| = 2|G|$.

Apply Lemma 2.1 with $F = E(X[V])$ and $\ell = 6$. As a result, there exists a 6-circuit C' such that $|F \cap E(C')| \geq 4$ (notice that, $|F \cap E(C)| = 2$ for C given in (2)). Thus C' lies entirely in V, or shares exactly one vertex with U.

In the former case $|a| = 6$, $|G| \in \{6, 18\}$, and $|V(X)| \in \{8, 24\}$ (recall that $|b| = |G_U| = 3$). By [4, Table], $X \cong Q_3$ or $GP(12, 5)$. Each of these graphs, however, is impossible. In fact, Q_3 has girth 4, and $GP(12, 5)$ has no abelian automorphism group with two orbits on the vertex set and in the same time acting unfaithfully on one of these orbits (this we found by the help of the computer package MAGMA [2]).

In the latter case C' can be written as

$$C' : u \ \sim \ v \ \sim \ va \ \sim \ va^2 \ \sim \ va^3 \ \sim \ va^4 \ \sim \ u,$$

where $u \in U$. Since $u \sim v$, $ua^4 \sim va^4$ also holds. As va^4 has exactly one neighbor in U, $ua^4 = u$, and so $a^4 \in G_U$. Recall that the order $|a| \geq 5$. Since also $|G_U| = 3$ and $G = \langle G_U, a\rangle$, we finally get $G = \langle a\rangle, |G| = 12$, and so $X \cong GP(8,3)$. We compute by MAGMA that this is indeed possible, $G \cong \mathbb{Z}_{12}$, and it has two orbits of lengths 4 and 12 respectively.

Case 2. $|G_U| = 2$.

In this case $|V| = |G|$, $|U| = \frac{1}{2}|G|$, $|V[X]| = \frac{3}{2}|G|$, and $|E(X)| = \frac{9}{4}|G|$. In particular, $|G|$ is divisible by 4. Notice that, $X[U]$ consists of a 1-factor, and G acts transitively, hence regularly, on the edges of $X[U]$. Let $H \leq G$ be the subgroup defined by $H = \{g \in G : eg = g \text{ for every } e \in E(X[U])\}$. It follows that $H \cong \mathbb{Z}_2^2$. It is clear that $G_U < H$. The connectedness of X implies that $\langle H, a\rangle$ is transitive on V, and hence $G = \langle H, a\rangle$.

Apply Lemma 2.1 with $F = E(X) \setminus E(X[V]) \setminus E(X[U])$ and $\ell = 6$. As a result, there exists a 6-circuit C' such that $|F \cap E(C')| \leq 2$. There are three possibilities for C': it lies entirely in V, or shares exactly one vertex with U, or shares exactly two adjacent vertices with U.

In the first case $|a| = 6$, and thus $|G| = |H| \cdot |\langle a\rangle|/|H \cap \langle a\rangle|$. Hence $|G| \in \{12, 24\}$, $|V(X)| \in \{18, 36\}$. By [4, Table], $X \cong$ F18 (the Pappus graph). This case we exclude by MAGMA, F18 has no abelian automorphism group with two orbits on the vertex set and in the same time acting unfaithfully on one of these orbits.

In the second case C' can be written as

$$C' : u \ \sim \ v \ \sim \ va \ \sim \ va^2 \ \sim \ va^3 \ \sim \ va^4 \ \sim \ u,$$

where $u \in U$. Reasoning as in the previous case, we get that $a^4 \in G_U$. Since $|G_U| = 2$ and $|a| \geq 5$, $|a| = 8$ and $H \cap \langle a\rangle = G_U$. Then $|G| = 16$, $|V(X)| = 24$, and so $X \cong GP(12,5)$. This graph was already excluded.

Finally, in the third case C' can be written as

$$C' : u_1 \ \sim \ u_2 \ \sim \ v \ \sim \ va \ \sim \ va^2 \ \sim \ va^3 \ \sim \ u_1,$$

where $u_1, u_2 \in U$. Then $u_2a^3 \sim va^3$, and hence $u_2a^3 = u_1$. Thus the edge $\{u_1a^3, u_2a^3\}$ must be equal to the edge $\{u_1, u_2\}$, showing that $u_1a^3 = u_2$, and $a^3 \in H$. As $|a| \geq 5$ and $H \cong \mathbb{Z}_2^2$, we see that $|a| = 6$, $|G| = 12$, $|V(X)| = 18$, and so $X \cong$ F18. This was already excluded. □

3 Bipartite graphs

In this section we turn to the case when G is faithful on both U and V. Till the end of Section 4 let us fix vertices $u \in U$ and $v \in V$ such that $\{u, v\} \in E(X)$. Then

$$U = \{ug : g \in G\} \text{ and } V = \{vg : g \in G\}. \tag{3}$$

Also, $ug \neq ug'$ and $vg \neq vg'$ whenever $g \neq g'$.

The goal of this section is to derive the following theorem:

Theorem 3.1. *Suppose that* G *acts faithfully on both* U *and* V, *and* X *is a bipartite graph with color classes* U *and* V. *Then* $(X, G) \cong (Q_3, \mathbb{Z}_4)$ *or* $(GP(8,3), \mathbb{Z}_8)$ *or* $X \cong H(G; a, b)$ *and* $G = \langle a, b \rangle$, *where* $a^\sigma = ba^{-1}$ *and* $b^\sigma = a^{-1}$ *for some* $\sigma \in \mathrm{Aut}(G)$.

The proof of Theorem 3.1 will be given in the end of the section. For this section X will be assumed to be bipartite with color classes U and V. Then there exist $a, b \in G$ suh that

$$E(X) = \{ \ \{ug, vg\}, \{ug, vga\}, \{ug, vgb\} : g \in G \ \}. \tag{4}$$

Then $G = \langle a, b \rangle$ by the connectedness of X. In fact, X is isomorphic to the Haar graph $H(G; a, b)$ defined in Example 1.2.

Let $N \leq G$. The orbit of $ug \in U$ ($vg \in V$, respectively) under N will be written as uNg (vNg, respectively). Suppose, in addition, that $N \neq G$ and $N \lhd \mathrm{Aut}(X)$. Then the orbits of N (N-orbits for short) form a *system of imprimitivity* for the permutation group $\mathrm{Aut}(X)$. Let K be the kernel of the action of $\mathrm{Aut}(X)$ on the set of orbits of N, i.e., $K = \{g \in \mathrm{Aut}(X) : Og = O$ for every N-orbit $O\}$. It is clear that $N \leq K$, and that K has the same orbits as N. For a vertex w, denote by K_w the stabilizer of w in K. Using that X is cubic and $N \neq G$, we obtain that if $w' \sim w$, then w and w' belong to different N-orbits. This implies that K_w fixes every neighbor of w. Combining this with the connectedness of X it follows in turn that, $|K_w| = 1$, $|K| \leq |N|$, and so $K = N$.

For $g \in \mathrm{Aut}(X)$, let $\overline{g}$ denote the permutation of the set of N-orbits induced by g, and for a subgroup $H \leq \mathrm{Aut}(X)$, let $\overline{H} = \{\overline{g} : g \in H\}$. Then $\overline{H} \cong H/(H \cap K)$, and since $K = N$, the following isomorphisms holds:

$$\overline{H} \cong H/(H \cap N) \cong HN/N. \tag{5}$$

In the sequel we shall consider the *factor graph* X/N defined as follows: the vertices are the N-orbits, i.e.,

$$V(X/N) = \{uNg : g \in G\} \ \cup \ \{vNg : g \in G\}; \tag{6}$$

and two orbits O and O' are adjacent if and only if there exist $w \in O$ and $w' \in O'$ such that $w \sim w'$ in X. It is not hard to see that X/N is a connected and cubic graph, the elements in $\overline{\mathrm{Aut}(X)}$ act as automorphisms of X/N, and $\overline{\mathrm{Aut}(X)}$ is transitive on $E(X/N)$ (compare these also with the previous paragraph). Obviously, $\overline{G} \leq \overline{\mathrm{Aut}(X)}$ and (5) gives that $\overline{G} \cong GN/N = G/N$. In what follows we write G/N for the permutation group $\overline{G}$.

In the next lemma we choose N to be the *core* of G in $\mathrm{Aut}(X)$, i.e., N is the largest normal subgroup of $\mathrm{Aut}(X)$ contained in G.

Lemma 3.2. *Let* N *be the core of* G *in* $\mathrm{Aut}(X)$ *and suppose that* $N \neq G$. *Then* $(X/N, G/N) \cong (K_{3,3}, \mathbb{Z}_3)$ *or* $(Q_3, \mathbb{Z}_4)$ *or* $(\mathrm{F14}, \mathbb{Z}_7)$.

Proof. Let $M \leq G$ such that $\overline{M} \trianglelefteq \mathrm{Aut}(X/N)$. Since $\overline{\mathrm{Aut}(X)} \leq \mathrm{Aut}(X/N)$, $\overline{M} \trianglelefteq \overline{\mathrm{Aut}(X)}$, and therefore, $MN/N \trianglelefteq \mathrm{Aut}(X)/N$, see (5). This gives that $MN \trianglelefteq \mathrm{Aut}(X)$. As $MN \leq G$ and N is the core of G in $\mathrm{Aut}(X)$, $MN = N$, and $\overline{M}$ is trivial. We conclude that $\overline{G} = G/N$ has trivial core in $\mathrm{Aut}(X/N)$. Using also that G/N is abelian with two orbits on $V(X/N)$ given in (6), it is sufficient to prove Lemma 3.2 for the particular case when N is trivial.

By Tutte's Theorem (see the introduction), X is s-regular for some $s \leq 5$. Thus $|\mathrm{Aut}(X)| = |V(X)| \cdot 3 \cdot 2^{s-1} = |G| \cdot 3 \cdot 2^s$, and hence $|\mathrm{Aut}(X) : G| = 3 \cdot 2^s$. Let us consider the action of $\mathrm{Aut}(X)$ on the set of right G-cosets in $\mathrm{Aut}(X)$. Then G acts as a point stabilizer, and as the core N is trivial, this action is faithful. Equivalently, G is embedded into $S_{3 \cdot 2^s - 1}$, and we write $G \leq S_{3 \cdot 2^s - 1}$. It was proved in [3, Theorem 1] that, if $n = 3m + 2$ and $A \leq S_n$ is abelian, then

$$|A| \leq 2 \cdot 3^m. \tag{7}$$

Moreover, equality holds above if and only if $A \cong \mathbb{Z}_2 \times \mathbb{Z}_3^m$.

Let $s = 1$. Then $G \leq S_5$, and thus $|G| \leq 6$. Then $|V(X)| \leq 12$, however, no graph in [4, Table] with less than 13 vertices is 1-regular. Observe next that X has the 6-circuit:

$$C : v \sim u \sim va \sim uab^{-1} \sim vab^{-1} \sim ub^{-1} \sim v.$$

Since X is bipartite, its girth is 4 or 6. If the girth is 4, then $(X, G) \cong (K_{3,3}, \mathbb{Z}_3)$ or $(Q_3, \mathbb{Z}_4)$, see Table 1. For the rest of the proof let the girth be equal to 6.

Let $s = 2$. Then $G \leq S_{11}$, $G = \langle a, b\rangle$, where a, b are defined in (4), and hence $|G| < 54$ follows from (7) and the remark afterward. Then $|V(X)| < 108$, and by [4, Table], $|V(X)| \in \{16, 24, 32, 50, 54, 72, 96, 104\}$. Each of these numbers can be excluded by the help of MAGMA.

Finally, let $s > 2$. Since X has girth 6, it follows by [5, Theorem 2.3] that, $X \cong$ F14 or F18 or $GP(10, 3)$. A direct computer check shows that $(X, G) \cong (\mathrm{F14}, \mathbb{Z}_7)$. $\square$

In our next proof we make use of properties of voltage graphs. We review next the relevant definitions and notations (for more details, we refer to [13]). Let Y be a finite simple graph and K be a finite group. For an arc $x = (w, w') \in A(Y)$ we set $x^{-1} = (w', w)$. A *K-voltage assignment* of Y is a mapping $\zeta : A(Y) \to K$ with the property $\zeta(x^{-1}) = \zeta(x)^{-1}$ for every $x \in A(Y)$. The values of ζ are called *voltages* and K is called the *voltage group*. Voltages are naturally extended to directed walks $W : w_1 \sim w_2 \sim \cdots \sim w_n$ by letting

$$\zeta(W) = \prod_{i=1}^{n-1} \zeta((w_i, w_{i+1})).$$

Fix a spanning tree T of Y. Then every edge not in $E(T)$ together with the edges in $E(T)$ span a unique circuit of Y, and we shall refer to the circuits obtained in this manner as the *base circuits of Y relative to T*. The K-voltage assignment ζ is called *T-reduced* if $\zeta(x) = 1_K$ whenever x is an arc in T.

The *voltage graph* $Y \times_\zeta K$ is defined to have vertex set $V(Y) \times K$, and edge set

$$E(Y \times_\zeta K) = \Big\{\{(w, k), (w', \zeta(x)k)\} : x = (w, w') \in A(Y) \text{ and } k \in K\Big\}. \qquad (8)$$

The voltage group K induces an automorphism group of $Y \times_\zeta K$ through the action $k_{right} : (w, l) \mapsto (w, lk)$, $w \in V(Y)$ and $k, l \in K$. We set $K_{right} = \{k_{right} : k \in K\}$. Let $g \in \mathrm{Aut}(Y \times_\zeta K)$ such that it normalizes K_{right}. This implies that, if $(w, k) \in V(Y \times_\zeta K)$ and $(w, k)g = (w', k')$, then w' does not depend on the choice of $k \in K$, and the mapping $w \mapsto w'$ is a well-defined permutation of $V(Y)$. The latter permutation will be called the *projection* of g, obviously, it belongs to $\mathrm{Aut}(Y)$. On the other hand, an automorphism of $\mathrm{Aut}(Y)$ is said to *lift* to an automorphism of $Y \times_\zeta K$ if it is the projection of some automorphism of $Y \times_\zeta K$. The following "lifting lemma" is a special case of [16, Theorem 4.2]:

Theorem 3.3. *Let $Y \times_\zeta K$ be a connected voltage graph, where K is an abelian group, and ζ is a T-reduced K-voltage assignment. Then $\sigma \in \mathrm{Aut}(Y)$ lifts to an automorphism of $Y \times_\zeta K$ if and only if there exists some $\sigma_* \in \mathrm{Aut}(K)$ such that*

$$\sigma_*(\zeta(C)) = \zeta(C\sigma)$$

for every directed base circuit C relative to T.

Lemma 3.4. *Let N be the core of G in $\mathrm{Aut}(X)$ and suppose that $N \neq G$, N is nontrivial such that $(X/N, G/N) \cong (Q_3, \mathbb{Z}_4)$. Then $(X, G) \cong (GP(8, 3), \mathbb{Z}_8)$.*

Proof. Recall that $G = \langle a, b\rangle$, where a and b are defined in (4). We may assume that Na is a generator of G/N. It is not hard to see that $Nb \neq N$ and $Nb \neq Na$, hence $Nb = Na^t$ for some $t \in \{2, 3\}$. Also, there exist $n_1, n_2 \in N$ such that $a^4 = n_1$ and $b = n_2a^t$. Since $G = \langle a, b\rangle$, $N = \langle n_1, n_2\rangle$. Here we consider only the case that $t = 2$ (the case that $t = 3$ can be treated in the same manner).

We proceed by defining a N-voltage assignment of the factor graph X/N (in fact, the cube Q_3). By (6),

$$V(X/N) = \{uNa^i : i \in \{0, 1, 2, 3\}\} \cup \{vNa^i : i \in \{0, 1, 2, 3\}\}.$$

We depict X/N in Figure 1, where also specify a spanning tree T by the dotted edges, and a T-reduced N-voltage assignment $\zeta : A(X/N) \to N$. We claim that $X/N \times_\zeta N \cong X$. For an easier notation, set $\widehat{X} = X/N \times_\zeta N$. Then $V(\widehat{X}) = V(X/N) \times N$. In fact, we are going to show that the mapping $f : V(\widehat{X}) \to V(X)$, defined by

$$f\colon (uNa^i, n) \mapsto una^i \text{ and } (vNa^i, n) \mapsto vna^i,$$

is an isomorphism from $\widehat{X}$ to X.

Let $g \in G$. Then g factors as $g = ma^j, m \in N, j \in \{0, 1, 2, 3\}$, define the permutation $\widehat{g}$ of $V(\widehat{X})$ by

$$\widehat{g}\colon (uNa^i, n) \mapsto \begin{cases} (uNa^{i+j}, mn) & \text{if } i + j < 4 \\ (uNa^{i+j-4}, mnn_1) & \text{if } i + j \geq 4 \end{cases}$$



Figure 1: Voltage assignment ζ of X/N.

and

$$\widehat{g}\colon (vNa^i, n) \mapsto \begin{cases} (vNa^{i+j}, mn) & \text{if } i+j<4 \\ (vNa^{i+j-4}, mnn_1) & \text{if } i+j \geq 4. \end{cases}$$

Notice that, actually, $\widehat{g} = fgf^{-1}$ holds for every $g \in G$. This gives that $\widehat{G} = fGf^{-1}$. We are going to show at first that $\widehat{g}$ is an automorphism of $\widehat{X}$. The statement follows quickly if $g = n \in N$ because then $\widehat{n} = n_{right}$. Since $G = \langle N, a\rangle$, it is enough to show that $\widehat{a} \in \mathrm{Aut}(\widehat{X})$. Let $\widehat{w}_1$ and $\widehat{w}_2$ be two adjacent vertices of $\widehat{X}$. By definition (8), we may write $\widehat{w}_1 = (uNa^{i_1}, n)$ and $\widehat{w}_2 = (vNa^{i_2}, \zeta(x)n)$, where $x = (uNa^{i_1}, vNa^{i_2})$ is an arc of X/N, and ζ is defined in Figure 1. We derive below that $\widehat{w}_1\widehat{a} \sim \widehat{w}_2\widehat{a}$:

Let $i_1 < 3$ and $i_2 < 3$, or $i_1 = i_2 = 3$. Then $\zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big) = \zeta(x)$. Set $\varepsilon = 0$ if $i_1 < 3$ and $i_2 < 3$, and $\varepsilon = 1$ if $i_1 = i_2 = 3$. Then the statement follows as

$$\begin{aligned} \widehat{w}_1\widehat{a} &= (uNa^{i_1+1}, nn_1^{\varepsilon}) \sim \Big(vNa^{i_2+1}, \zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big)nn_1^{\varepsilon}\Big) \\ &= (vNa^{i_2+1}, \zeta(x)nn_1^{\varepsilon}) = \widehat{w}_2\widehat{a}. \end{aligned}$$

If $i_1 < 3$ and $i_2 = 3$, then $\zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big) = n_1\zeta(x)$, and

$$\begin{aligned} \widehat{w}_1\widehat{a} &= (uNa^{i_1+1}, n) \sim \Big(vNa^{i_2+1}, \zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big)n\Big) \\ &= (vNa^{i_2+1}, \zeta(x)nn_1) = \widehat{w}_2\widehat{a}. \end{aligned}$$

Finally, if $i_1 = 3$ and $i_2 < 3$, then $\zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big) = n_1^{-1}\zeta(x)$, and

$$\begin{aligned} \widehat{w}_1\widehat{a} &= (uNa^{i_1+1}, nn_1) \sim \Big(vNa^{i_2+1}, \zeta\big((uNa^{i_1+1}, vNa^{i_2+1})\big)nn_1\Big) \\ &= (vNa^{i_2+1}, \zeta(x)n) = \widehat{w}_2\widehat{a}. \end{aligned}$$

Set $\widehat{G} = \{\widehat{g} : g \in G\} \leq \mathrm{Aut}(\widehat{X})$. It can be seen that $\widehat{G}$ has two orbits on $V(\widehat{X})$ induced by the vertices $(uN, 1_N)$ and $(vN, 1_N)$ respectively. Let $N_{\widehat{X}}((uN, 1_N))$ denote the neighborhood of $(uN, 1_N)$ in $\widehat{X}$. Then

$$N_{\widehat{X}}((uN, 1_N)) = \{(vN, 1_N), (vNa, 1_N), (vNa^2, n_2)\}.$$

The above set is mapped by f to the set $\{v, va, vn_2a^2\}$, and since $b = n_2a^2$, this is equal to the neighborhood of u in the graph X. Also, as f maps $(uN, 1_N)$ to u, we obtain that, for any edge $e \in E(\widehat{X})$ such that e is incident with $(uN, 1_N)$, ef will be an edge of X (here ef denotes the image of e under the mapping f). We derive that f is an isomorphism by showing that the same holds for any edge $e \in E(\widehat{X})$. Observe that, there exists a suitable $\widehat{g} \in \widehat{G}$ so that $e\widehat{g}$ is incident with $(Nu, 1_N)$, put $e' = e\widehat{g}$. Then $ef = (e'\widehat{g}^{-1})f = (e'(fg^{-1}f^{-1}))f = (e'f)g^{-1}$, showing that ef is indeed an edge of X.

Since $N \trianglelefteq \mathrm{Aut}(X)$, $fNf^{-1} \trianglelefteq f\,\mathrm{Aut}(X)f^{-1} = \mathrm{Aut}(\widehat{X})$. As $fNf^{-1} = \widehat{N} = N_{right}$, we see that every $g \in \mathrm{Aut}(\widehat{X})$ has a projection $\tilde{g}$. Let us denote by P the projection of $\mathrm{Aut}(\widehat{X})$, i.e., $P = \{\tilde{g} : g \in \mathrm{Aut}(\widehat{X})\}$. In particular, the projection of $\widehat{G}$ is generated by the 4-cycle $\alpha = (uN\ uNa\ uNa^2\ uNa^3)(vN\ vNa\ vNa^2\ vNa^3)$, hence $\alpha \in P$. Using also that P is transitive on $E(X/N)$, we compute by Magma that $P = \mathrm{Aut}(X/N)$. Thus for the automorphism $\beta = (uNa^2\ uNa^3)(vNa\ vNa^2)$ of X/N, β lifts to an automorphism of $\widehat{X}$.

We apply Theorem 3.3 to $\widehat{X} = X/N \times_\zeta N$ with $\sigma = \beta$ and the following directed base circuits relative to T:

$$C : uN \ \sim\ vNa^2 \ \sim\ uNa \ \sim\ vNa \ \sim\ uN,$$

$$C' : uNa \ \sim\ vNa^3 \ \sim\ uNa^2 \ \sim\ vNa^2 \ \sim\ uNa,$$

$$C'' : uN \ \sim\ vNa \ \sim\ uNa \ \sim\ vNa^2 \ \sim\ uNa^2 \ \sim\ vN \ \sim\ uN.$$

Let σ_* be the automorphism of N given in Theorem 3.3. Since $\zeta(C) = \zeta(C') = n_2$, $\zeta(C\beta) = \sigma_*(n_2) = \zeta(C'\beta)$. This gives $n_2^{-1} = n_1n_2^2$. Also, $\zeta(C'') = n_1n_2$ and $\zeta(C''\beta) = 1_N$, hence $\sigma_*(n_1n_2) = 1_N$, and so $n_1n_2 = 1_N$. Finally, $n_1 = n_2$, $n_1^2 = 1_N$, and so $|N| = 2$. Thus $X \cong GP(8,3)$, and we compute by Magma that $G \cong \mathbb{Z}_8$. $\square$

Proof of Theorem 3.1. As noted before, $X \cong H(G; a, b)$, where $a, b \in G$ are defined in (4). We are going show the following property: if $X \ncong Q_3$ nor $GP(8,3)$, then there exists a subgroup $H \leq \mathrm{Aut}(X)$ such that

$$H \text{ is transitive on } E(X) \text{ and } G \trianglelefteq H. \tag{9}$$

Let N be the core of G in $\mathrm{Aut}(X)$. If $N = G$, then (9) holds by letting $H = \mathrm{Aut}(X)$. Let $N < G$. By Lemmas 3.2 and 3.4, $(X/N, G/N) \cong (K_{3,3}, \mathbb{Z}_3)$ or $(\mathrm{F}14, \mathbb{Z}_7)$. Using that $\overline{\mathrm{Aut}(X)}$ is transitive on $E(X/N)$, we obtain by Magma that in either case $\overline{\mathrm{Aut}(X)}$ contains a subgroup K such that K is transitive on $E(X/N)$ and $\overline{G} \trianglelefteq K$. Now, (9) holds by letting H to be the subgroup $H \leq \mathrm{Aut}(X)$ for which $G \leq H$ and $\overline{H} = K$.

Let H_u denote the stabilizer of u in H. The orbit of v under H_u is equal to $\{v, va, vb\}$, thus there exists an element $\alpha \in H_u$ which acts on this orbit as a 3-cycle. We may write that $v\alpha = va$, $(va)\alpha = vb$ and $(vb)\alpha = v$. As α normalizes G and sends u to u and v to va, there exists an automorphism $\sigma \in \mathrm{Aut}(G)$ such that α acts on $V(X)$ as

$$\alpha : ug \mapsto ug^\sigma \text{ and } vg \mapsto v(g^\sigma a).$$

In particular, $vb = (va)\alpha = v(a^\sigma a)$ and $v = (vb)\alpha = v(b^\sigma a)$. Thus $b = a^\sigma a$ and $1_G = b^\sigma a$, hence a and b satisfy the conditions given in Theorem 3.1, and so X is symmetric (see Example 1.2). $\square$

4 Proof of part (B)

Another partial case of Theorem 1.3.(B) follows directly from [14, Theorem 1.1]:

Theorem 4.1. *Suppose that G acts faithfully on both U and V, any vertex in U has one neighbor in V, and any vertex in V has one neighbor in U. Then (X, G) is isomorphic to one of the following pairs:*

$$(F40, \mathbb{Z}_2 \times \mathbb{Z}_{10}),\ (GP(4,1), \mathbb{Z}_4),\ (GP(4,1), \mathbb{Z}_2^2),\ (GP(5,2), \mathbb{Z}_5),\ (GP(8,3), \mathbb{Z}_8),$$
$$(GP(10,2), \mathbb{Z}_{10}),\ (GP(10,3), \mathbb{Z}_{10}),\ (GP(12,5), \mathbb{Z}_{12}),\ (GP(24,5), \mathbb{Z}_{24}).$$

In view of Theorems 3.1 and 4.1, Theorem 1.3.(B) reduces to the following theorem, which we are going to prove in the end of the section:

Theorem 4.2. *Suppose that G acts faithfully on both U and V, any vertex in U has two neighbors in V, and any vertex in V has two neighbors in U. Then (X, G) is isomorphic to one of the following pairs:*

$$(K_4, \mathbb{Z}_2),\ (Q_3, \mathbb{Z}_2^2),\ (GP(8,3), \mathbb{Z}_2 \times \mathbb{Z}_4),\ (GP(12,5), \mathbb{Z}_2 \times \mathbb{Z}_6).$$

The orbits U and V are as defined in (3), and as for the edges, there exist $a, b, c \in G$ such that

$$E(X) = \big\{ \{ug, uga\}, \{vg, vgb\}, \{ug, vg\}, \{ug, vgc\} : g \in G \big\}. \tag{10}$$

Clearly, a and b are involutions, and it can be easily shown that $G = \langle a, b, c\rangle$.

Lemma 4.3. *Let N be the core of G in* Aut(X). *Then $(X/N, G/N) \cong (K_4, \mathbb{Z}_2)$ or $(Q_3, \mathbb{Z}_2^2)$.*

Proof. Similarly to the proof of Lemma 3.2, it is sufficient to prove the lemma for the case when N is trivial. If X has girth at most 4, then $(X, G) \cong (K_4, \mathbb{Z}_2)$ or $(Q_3, \mathbb{Z}_2^2)$, see Table 1. Let the girth of X be larger than 4. Then $a \neq b$, for otherwise, we have the 4-circuit: $u \sim ua \sim va = vb \sim v \sim u$. If $\langle a, b\rangle \cap \langle c\rangle$ is trivial, then

$$G = \langle a\rangle \times \langle b\rangle \times \langle c\rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_n. \tag{11}$$

Otherwise, c must be of even order, say $2n$, and $c^n \in \langle a, b\rangle$. We may assume without loss of generality that $c^n \neq a$, and thus

$$G = \langle a\rangle \times \langle c\rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_{2n},\ \text{and } b = c^n \text{ or } ac^n. \tag{12}$$

By Tutte's Theorem X is s-regular for some $s \leq 5$. Reasoning as in the proof of Lemma 3.2, we see that G is embedded into $S_{3 \cdot 2^s - 1}$, and we write $G \leq S_{3 \cdot 2^s - 1}$.

If $s = 1$, then $\mathbb{Z}_2 \times \mathbb{Z}_2 \leq G \leq S_5$, from which it follows at once that $|G| = 4$, and $X \cong Q_3$, which is excluded as the girth of X is larger than 4.

Let $s = 2$. Then $G \leq S_{11}$, and thus $n \leq 13$ by (7). A direct computer check shows that if G is given as in (11) and $n \leq 13$, then X is not edge-transitive. Furthermore, if G is given as in (12) and $n \leq 13$, then X is edge-transitive in two cases: $n = 2$ or 3, and $b = ac^n$. The corresponding graphs are $GP(8,3)$ and $GP(12,5)$ respectively, however, then the core of G in the full automorphism group has order 2 and 3 respectively.

Let $s \geq 3$. By the previous paragraph we may assume that $n > 13$. The graph X has the 8-circuit:

$$u \;\sim\; ua \;\sim\; va \;\sim\; vab \;\sim\; uab \;\sim\; ub \;\sim\; vb \;\sim\; v \;\sim\; u.$$

Thus X must have an 8-circuit starting with the 3-arc $u \sim vc \sim uc \sim vc^2$, too. Let us write such an 8-circuit in the form:

$$C : u \sim vc \sim uc \sim vc^2 \sim xc^2\alpha \sim x'c^2\alpha\beta \sim x''c^2\alpha\beta\gamma \sim x'''c^2\alpha\beta\gamma\delta \sim u,$$

where $x, x', x'', x'''' \in \{u, v\}$ and $\alpha, \beta, \gamma, \delta \in \{1_G, a, b, c, c^{-1}\}$. Put $\eta = c^2\alpha\beta\gamma\delta$. Then

$$\eta = c^i a^j b^k \text{ for some } i \in \{0, \ldots, 4\} \text{ and } j, k \in \{0, 1\}.$$

Since $u \sim x'''\eta$ and $x''''\eta \neq vc$, $\eta \in \{1_G, a\}$. Thus $1_G = \eta^2 = c^{2i}a^{2j}b^{2k} = c^{2i}$. This implies that $i = 0$, for otherwise, the order of c is at most 8, hence $n \leq 8$ (see (11) and (12)), which contradicts the assumption that $n > 13$. It is easy to see that $i = 0$ happens only in one case:

$$C : u \sim vc \sim uc \sim vc^2 \sim vc^2b \sim ucb \sim vcb \sim ub \sim u.$$

Then, however, $\eta = b$, contradicting that $\eta \in \{1_G, a\}$ and $a \neq b$. $\square$

Proof of Theorem 4.2. Let N be the core of G in $\mathrm{Aut}(X)$. If N is trivial, then X and G are determined in Lemma 4.3. For the rest of the proof let N be non-trivial. We proceed by the two cases of Lemma 4.3.

Case 1. $X/N \cong K_4$ and $G/N \cong \mathbb{Z}_2$.

It is easily seen that none of a, b and c is in N, and hence $G = N \times \langle a \rangle$, $b = n_1 a$, and $c = n_2 a$ for some $n_1, n_2 \in N$. Since $G = \langle a, b, c \rangle$, $N = \langle n_1, n_2 \rangle$. Also, since $a \neq b$ are involutions, n_1 is an involution.

Assume for the moment that N is not a 2-group, and let p be an odd prime divisor of $|N|$. Then the subgroup $M = \langle n_1, n_2^p \rangle$ is the unique subgroup in N of index p, hence it is characteristic in N. Using also that $N \trianglelefteq \mathrm{Aut}(X)$, we obtain that $M \trianglelefteq \mathrm{Aut}(X)$. The quotient graph X/M is a connected cubic symmetric graph on $4p$ points. Also, G/M is an abelian subgroup of $\mathrm{Aut}(X/M)$ with two orbits such that there are two matchings between these orbits. Since X has $4p$ points, it was proved in [9, Theorem 6.2] that $X \cong Q_3$ or $GP(8,3)$ or $GP(10,3)$ or $GP(10,5)$ or $F28$ (the Coxeter graph). However,



Figure 2: Voltage assignment ζ of X/N.

we compute by Magma that none of these graphs have an abelian automorphism group with two orbits of equal lengths and in the same time there are two matchings between the orbits, and therefore, N must be a 2-group.

Recall that $N = \langle n_1, n_2 \rangle$ and n_1 is involution. Let $|N| = 2^t$, and assume for the moment that $|N| \geq 8$. Then there is a characteristic subgroup $M \leq N$ with $|N : M| = 8$. This can be seen at once if $|N| = 8$ or N is cyclic; while when none of these holds, then $N \cong \mathbb{Z}_2 \times \mathbb{Z}_{2^{t-1}}$ for $t \geq 4$, and M is the unique subgroup isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_{2^{t-3}}$. Since M is characteristic in N and $N \trianglelefteq \mathrm{Aut}(X)$, $M \trianglelefteq \mathrm{Aut}(X)$, and the quotient graph X/M is a connected cubic symmetric graph on 32 points. By [4, Table], $X/M \cong$ F32 (the Dyck graph). This graph, however, has no abelian automorphism group with two orbits of equal lengths and in the same time there are two matchings between the orbits. We conclude that $|N| = 2$ or $N = |4|$. If $|N| = 2$, then $X \cong Q_3$, and $G \cong \mathbb{Z}_2^2$, see Table 1. If $|N| = 4$, then $X \cong GP(8,3)$, and we compute by Magma that $G \cong \mathbb{Z}_2 \times \mathbb{Z}_4$, and $N \cong \mathbb{Z}_4$.

Case 2. $X/N \cong Q_3$ and $G/N \cong \mathbb{Z}_2^2$.

Again, none of a, b, c is in N. Assume for the moment that $Na = Nb$. Since $G = \langle a, b, c \rangle$, $G/N = \langle Na, Nc \rangle$. Let us consider the quotient graph $X/N \cong Q_3$. As N is the core of G in $\mathrm{Aut}(X)$, G/N has trivial core in $\mathrm{Aut}(X/N)$. By (6), $V(X/N) = \{uN, uNa, uNc, uNac, vN, vNa, vNc, vNac\}$. Now, uN has neighbors as uNa, vN and vNc, vertices at distance 2 as $vNa, vNac, uNc$, and so its antipodal vertex is $uNac$. Similar considerations yield that the antipodal pairs of X/N are the orbits of the involution Nac, hence Nac is the central involution in $\mathrm{Aut}(X/N)$. This, however, contradicts that G/N has trivial core in $\mathrm{Aut}(X/N)$. We conclude that $Na \neq Nb$.

Then $G/N = \langle Na, Nb \rangle$, $c = n_1 ab$ for some $n_1 \in N$, and finally $N = \langle n_1 \rangle$. We depict X/N in Figure 2, where also fix a spanning tree T shown by the dotted edges, and a T-reduced N-voltage assignment $\zeta : A(X/N) \to N$. As before, we set $\widehat{X} = X/N \times_\zeta N$, and define the mapping $f : V(\widehat{X}) \to V(X)$ by

$$f\colon (uNx, n) \mapsto unx \text{ and } (vNx, n) \mapsto vnx,\ x \in \{1_G, a, b, ab\}.$$

Reasoning as in the proof of Lemma 3.4, we obtain that f is an isomorphism from $\widehat{X}$ to X, and also that $\mathrm{Aut}(\widehat{X})$ projects to a subgroup of $\mathrm{Aut}(X/N)$ which is transitive on $E(X/N)$. Thus for the automorphism $\alpha = (uNa\ vNab\ vN)\big(vNa\ uNab\ vNb)$ of X/N, α lifts to an automorphism of $\widehat{X}$.

We apply Theorem 3.3 to $\widehat{X} = X/N \times_\zeta N$ with $\sigma = \alpha$ and the following directed base circuits relative to T:

$$C : uN\ \sim\ vNab\ \sim\ vNa\ \sim uNa,$$

$$C' : uNb\ \sim\ vNa\ \sim\ vNab\ \sim uNab \sim uNb.$$

Let σ_* be the automorphism of N given in Theorem 3.3. Since $\zeta(C) = \zeta(C') = n_1$, $\zeta(C\alpha) = \sigma_*(n_1) = \zeta(C'\alpha)$, which gives $n_1^{-2} = n_1$. Thus $|N| = 3$, $|V(X)| = 24$, $X \cong GP(12, 5)$, and we compute by Magma that $G \cong \mathbb{Z}_2 \times \mathbb{Z}_6$ and $N \cong \mathbb{Z}_3$. $\square$

Acknowledgements

In this research both authors have been supported in part by "Javna agencija za raziskovalno dejavnost Republike Slovenije", program no. P1-0285.

References

[1] Y. G. Baik, Y. Q. Feng, H. S. Sim, and M. Y. Xu, On the normality of Cayley graphs of abelian groups, *Algebra Colloq.* **5** (1998), 227–234.

[2] W. Bosma, J. Cannon, and C. Playoust, The Magma Algebra System I: The User Language, *J. Symbolic Comput.* **24** (1997), 235–265.

[3] J. M. Burns, B. Goldsmith, Maximal order abelian subgroups of symmetric groups, *Bull. London Math. Soc.* **21** (1989), 70–72.

[4] M. D. E. Conder and P. Dobcsányi, Trivalent symmetric graphs on up to 768 vertices, *J. Combin. Math. & Combin. Comp.* **40** (2002), 41–63.

[5] M. D. E. Conder, R. Nedela, Symmetric cubic graphs of small girth, *J. Combin. Theory Ser. B* **97** (2007), 757–768.

[6] M. D. E. Conder, R. Nedela, A refined classification of symmetric cubic graphs, *J. Algebra* **322** (2009), 722–740.

[7] S. F. Du, J. Q. Feng, J. H. Kwak, . M. Y. Xu, Cubic Cayley graphs on dihedral groups, *Math. Anal. Appl.* **7** (2004), 224–234.

[8] D. Ž. Djoković and G. L. Miller, Regular groups of automorphisms of cubic graphs, *J. Combin. Theory Ser. B* **29** (1980), 195–230.

[9] Y. Q. Feng and J. H. Kwak, Cubic symmetric graphs of order a small number time a prime or a prime square, *J. Combin. Theory Ser. B* **97** (2007), 627–646.

[10] Y. Q. Feng, R. Nedela, Symmetric cubic graphs of girth at most 7, *Acta Univ. M. Belii Math.* **13** (2006), 33–55.

[11] R. Frucht, J. E. Graver, and M. E. Watkins, The groups of the generalized Petersen graphs, *Proc. Camb. Phil. Soc.*, **70** (1971), 211-218.

[12] M. Hladnik, D. Marušič, and T. Pisanski, Cyclic Haar graphs, *Discrete Math.* **244** (2002), 137–152.

[13] J. L. Gross, T. W. Tucker, *Topological graph theory,* Wiley-Interscience, New York, 1987.

[14] I. Kovács, A. Malnič, D. Marušič and Š. Miklavič, One-mathcing bi-Cayley graph over abelian groups, *Europ. J. Combin.* **30** (2009), 602–616.

[15] K. Kutnar, D. Marušič, A complete classification of cubic symmetric graphs of girth 6, *J. Combin. Theory Ser. B* **99** (2009), 162–184.

[16] A. Malnič, Group actions, coverings and lifts of automorphisms, *Discrete Math.* **182** (1998), 203–218.

[17] D. Marušič, T. Pisanski, Symmetries of hexagonal graphs on the torus, *Croat. Chemica Acta* **73** (2000), 969–981.

[18] R. C. Miller, The trivalent symmetric graphs of girth at most six, *J. Combin. Theory Ser. B* **10** (1971), 163–182.

[19] T. Pisanski, A classification of cubic bicirculants, *Discrete Math.* **307** (2007), 567–758.

[20] W. T. Tutte, A family of cubical graphs, *Proc. Cambr. Philosoph. Soc.* **43** (1947), 459–474.

[21] S. J. Xu, W. Jin, Q. Shi, J. J. Li, The BCI-property of the Bi-Cayley graphs, *J. Guangxi Norm. Univ.: Nat. Sci. Edition* **26** (2008), 33–36.

[22] J. Xu, M. Y. Xu, Arc-transitive Cayley graphs of valency at most four on abelian groups *Southeast Asian Bull. Math.* **25** (2001), 355–363.